

Pediatric Discharge Instructions

Dosing Chart: Tylenol™ (acetaminophen)

Children 2 months and younger should not get Tylenol™ unless advised by your doctor.

Child's weight in pounds today _____	Equivalent to this weight in kilograms	Dose of Tylenol:	Children's Tylenol Elixir: 160 mg/ 5 mL	Chewable Tablets: 80 mg/tablet	Chewable Tablets: 160 mg/ tablet	Adult Strength Tylenol Tablets: 325 mg/ tablet
Pounds	Kilograms					
5	2.3	32 mg	1 mL	----	----	----
6 to 8	2.7-3.6	40 mg	1.25 mL	----	----	----
9 to 11	4.1-5	64 mg	2 mL	----	----	----
12 to 14	5.5-6.4	80 mg	2.5 mL	1 tablet	½ tablet	----
15 to 17	6.8-7.7	96 mg	3 mL	1 tablet	½ tablet	----
18	8.2	122 mg	3.75 mL	1 ½ tablets	½ tablet	----
19 to 23	8.6-10.5	128 mg	4 mL	1 ½ tablets	½ tablet	----
24 to 28	10.9-12.7	160 mg	5 mL	2 tablets	1 tablet	----
29 to 32	13.2-14.5	192 mg	6 mL	2 tablets	1 tablet	----
33 to 35	15-15.9	224 mg	7 mL	2 ½ tablets	1 tablet	----
36 to 46	16.4-20.9	240 mg	8 mL	3 tablets	1 ½ tablets	----
47 to 58	21.4-26.4	320 mg	10 mL	4 tablets	2 tablets	1 tablet
59 to 70	26.8-31.8	400 mg	12 mL	5 tablets	2½ tablets	1 tablet
71 to 90	32.3-40.9	480 mg	15 mL	6 tablets	3 tablets	1 ½ tablets
90+	40.9	620 mg	10 mL	8 tablets	4 tablets	1 ½ tablets

Take Tylenol (Acetaminophen) every 4 hours for fever or pain.
MAXIMUM OF 5 DOSES IN 24 HOURS.

DO NOT GIVE with ANY other medications that contain acetaminophen -- Tylenol, APAP, FeverAll Cetafen, Tempra, Panadol, Q-Pap, some Triaminic products and many other generic and combination products. Look for acetaminophen on the label. Any questions? Ask a pharmacist.

Pediatric Discharge Instructions

Dosing Chart: Motrin™ (ibuprofen)

For children aged 6 months and older

Child's weight in pounds today _____	Equivalent to this weight in kilograms	Dose of Ibuprofen:	Infant Ibuprofen Drops: 50 mg/ 1.25 mL	Children's Ibuprofen Elixir: 100 mg/5 mL	Chewable Tablets: 50 mg/ tablet	Chewable Tablets: 100mg/ tablet	Adult Strength Ibuprofen Tablets: 200mg/ tablet
Pounds	Kilograms						
12 to 13	5.5-5.9	50 mg	1.25 mL	_____	_____	_____	_____
14 to 17	6.4-7.7	60 mg	1.5 mL	3 mL	_____	_____	_____
18 to 21	8.2-9.5	80 mg	2 mL	4 mL	_____	_____	_____
22 to 26	10-11.8	100 mg	2.5 mL	5 mL	2 tablets	1 tablet	_____
27 to 35	12.3-15.9	120 mg	3 mL	6 mL	2 tablets	1 tablet	_____
36 to 39	16.4-17.7	160 mg	4 mL	8 mL	3 tablets	1½ tablets	_____
40 to 43	18.2-19.5	180 mg	4.5 mL	9 mL	3 tablets	1½ tablets	_____
44 to 52	20-23.6	200 mg	5 mL	10 mL	4 tablets	2 tablets	1 tablet
53 to 57	24.1-25.9	240 mg	6 mL	12 mL	4 tablets	2 tablets	1 tablet
58 to 65	26.4-29.5	260 mg	6.5 mL	13 mL	5 tablets	2½ tablets	1 tablet
66 to 70	30-31.8	300 mg	7.5 mL	15 mL	6 tablets	3 tablets	1 tablet
71 to 87	32.3-39.5	320 mg	8 mL	16 mL	6 tablets	3 tablets	1 tablet
88 to 120	40-54.5	400 mg	10 mL	20 mL	8 tablets	4 tablets	2 tablets
120+	54.5	Adult Dose	_____	_____	_____	4 tablets	2 tablets

Take Ibuprofen every 6 to 8 hours for fever, pain or inflammation.
MAXIMUM OF 4 DOSES IN 24 HOURS.

DO NOT GIVE Ibuprofen to children under age 6 months. Avoid other NSAIDs like naproxen or toradol. **DO NOT GIVE** with other medications that contain ibuprofen such as Advil, Motrin IB, and many generic products. Look for ibuprofen on the label. Any questions? Ask a pharmacist.